arXiv:1501.05620v1 [physics.flu-dyn] 21 Jan 2015

January 2015

Vortex solutions of the generalized Beltrami flows to the incompressible Euler equations

Minoru Fujimoto[1], Kunihiko Uehara[2] and Shinichiro Yanase[3]

[1]*Seika Science Research Laboratory, Seika-cho, Kyoto 619-0237, Japan*
[2]*Department of Physics, Tezukayama University, Nara 631-8501, Japan*
[3]*Graduate School of Natural Science and Technology, Okayama University, Okayama 700-8530, Japan*

Abstract

As for the solutions of the generalized Beltrami flows to the incompressible Euler equations besides the solutions separating radius and axial components, there are only several solutions found as the Hill's vortex solutions. We will present a series of vortex solutions in this category for the generalized Beltrami flows to the incompressible Euler equations.

PACS number(s): 47.10.ad, 47.15.ki, 47.32.-y

It is a well-known fact for many years that the Hill's spherical vortex [4] is one of exact solutions to the incompressible Euler equations. A characteristic feature of the solution is that the vorticity only exists inside the sphere, where the vorticity $\boldsymbol{\omega}$ is referred by the velocity $\boldsymbol{u}$ as

$$\boldsymbol{\omega} = \boldsymbol{\nabla} \times \boldsymbol{u}. \tag{1}$$

From a point of view in the differential equations, the Hill's solution belongs to the solutions of the generalized Beltrami flows

$$\boldsymbol{\nabla} \times (\boldsymbol{\omega} \times \boldsymbol{u}) = 0 \tag{2}$$

to the incompressible Euler equations. One category [2, 7, 9, 10] is made of the solutions separating radius and axial components in the cylindrical coordinates where the solutions

are described by the Bessel and exponential functions, the other is of the solutions as the Hill's vortex solution which are non-separable as for radius and axial components where there are only several solutions found so far.

The incompressible Euler equations

$$\frac{\partial \boldsymbol{u}}{\partial t} + (\boldsymbol{u} \cdot \nabla)\boldsymbol{u} = -\frac{1}{\rho}\nabla p, \tag{3}$$

$$\nabla \cdot \boldsymbol{u} = 0, \tag{4}$$

where p is the pressure and ρ is the density, are the non-linear differential equations because of the existence of the second term of the left-hand side in (3). The generalized Beltrami condition (2) is a linearization condition for the equations. This is seen by taking the curl of (3) as

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \nabla \times (\boldsymbol{\omega} \times \boldsymbol{u}) = 0. \tag{5}$$

As for *the exact solutions* to the incompressible Euler equations in this category, namely non-separable solutions, the Agrawal's[1] [1], the Berker's [2], the O'Brien's [6] and the Wang's solution [8] are known besides the Hill's vortex solution. These solutions are all axisymmetric flows which demand the conditions

$$\boldsymbol{u}(\eta, \varphi, z) = (u_\eta, 0, u_z), \tag{6}$$

$$\boldsymbol{\omega}(\eta, \varphi, z) = (0, \omega_\varphi, 0), \tag{7}$$

where we take the cylindrical coordinate (η, φ, z). When we introduce the Stokes stream function $\psi(\eta, \varphi, z)$, which automatically satisfies the continuity condition (4), the equations (3) reduce to

$$\frac{\partial}{\partial \eta}\left(\frac{1}{\eta}\frac{\partial \psi}{\partial \eta}\right) + \frac{1}{\eta}\frac{\partial^2 \psi}{\partial z^2} = -\omega_\varphi. \tag{8}$$

Meanwhile Marris and Aswani [5] demonstrated that the only solution for these axisymmetric generalized Beltrami flows is that ω_φ is proportional to η. Then (8) becomes

$$\frac{\partial^2 \psi}{\partial \eta^2} - \frac{1}{\eta}\frac{\partial \psi}{\partial \eta} + \frac{\partial^2 \psi}{\partial z^2} = -\alpha \eta^2, \tag{9}$$

where α is constant. There is a description "A simple, but useful, set of solutions" [3] of (9), which is

$$\psi = a_1\eta^4 + a_2\eta^2 z^2 + a_3\eta^2 + a_4\eta^2 z + a_5(\eta^6 - 12\eta^4 z^2 + 8\eta^2 z^4), \tag{10}$$

[1] $\psi = a_2\eta^2 z^2$ in (10) is a special case of the Agrawal solution, but this is a separable case when other coefficient are all zero.

where $8a_1 + 2a_2 = -\alpha$, but a_i's are otherwize arbitrary. In order to genarate the right-hand side of (9), we have to put the term of η^4 or $\eta^2 z^2$ in ψ, where *the exact solutions* above are all contained the either term or the both terms. In other words, the third,the forth or the fifth term of the left-hand side in (10) where the each coefficient a_3,a_4 or a_5 is arbitray, satisfies that the left-hand side of (9) equals zero.

The extention of the solution space for (9) is to add higher power terms for η and z which satisfy the same condition as a_3, a_4 or a_5 term. We present the result, a series of higher power terms, as

$$\begin{aligned} \psi &= a_1\eta^4 + a_2\eta^2 z^2 + a_3\eta^2 + a_4\eta^2 z \\ &\quad + c_n \sum_{k=0}^{n-1} a_{n,k}\eta^{2n-2k} z^{2k}, \end{aligned} \tag{11}$$

where $n \geq 3$, $a_{n,0} = 1$ and $a_{n,k} = \dfrac{(-4)^k \Gamma(n)\Gamma(n+1)}{\Gamma(2k+1)\Gamma(n-k)\Gamma(n-k+1)}$.

When the parameters are taken appropriately as $a_1, a_3, c_3 \neq 0$ and other coefficients vanish in (11), we reproduce the Wang's toroidal vortex shown in Fig.1.



Figure 1: $a_1 = 10, a_3 = -1, c_3 = 1/8$. The streamsurfaces have $\psi = -0.005, -0.01, 0, 0.1, 0.2$ from the innermost to the outer side.

The examples for higher order solutions are shown in Fig.2 and Fig.3. Their stream functions are

$$\psi = a_1\eta^4 + a_3\eta^2 + c_4\left(\eta^8 - 24\eta^6 z^2 + 48\eta^4 z^4 - \frac{64}{5}\eta^2 z^6\right) \tag{12}$$

and

$$\psi = a_1\eta^4 + a_3\eta^2 + c_5\left(\eta^{10} - 40\eta^8 z^2 + 160\eta^6 z^4 - 128\eta^4 z^6 + \frac{128}{7}\eta^2 z^8\right) \tag{13}$$



Figure 2: $a_1 = 10, a_3 = -1, c_4 = -1/8$. The streamsurfaces have $\psi = -0.005, -0.01, 0, 0.1, 0.2$ from the innermost to the outer side.



Figure 3: $a_1 = 10, a_3 = -1, c_5 = 1/16$. The streamsurfaces have $\psi = -0.005, -0.01, 0, 0.1, 0.2$ from the innermost to the outer side.

respectively.

Although we have to tune the parameters for a vortex like in the figures to appear in each orders, the higher order of η and z solutions we take, we get the streamsurface of $\psi = 0$ in the shape of from an oval as the Wang's vortex to more cylinder-like boundaries by the same order values of parameters. The toroidal vorticity $\omega_\varphi = \alpha\eta$ in each case though, these vorticity solutions should be the solutions inside the streamsurface of $\psi = 0$. The reason is that the global vorticity often generates some disturbances in the streamsurfaces and this will be easily confirmed when we draw the figures in larger ranges of η and z.

So, the remaining work we have to do is that we pour the perfect fluid outside of the boundaries of $\psi = 0$ and connect these solutions to the outer solutions with no-vorticity like the Hill's solution.

References

[1] H.L. Agrawal, *A new exact solution of the equations of viscous motion with axial symmetry*, Q.J. Mech. Appl. Math. **10**, 42-44 (1957).

[2] R. Berker, *Intégration des équations du mouvement d'un fluide visqueux incompressible*, Encyclopedia of Physics, ed. S. Flügge, vol. VIII/2, Berlin: Springer-Verlag, 1-384 (1963).

[3] P. Drazin and N. Riley, page 51 in *"The Navier-Stokes Equations"*, London Mathematical Society Lecture Note Series 334, Cambridge Univ. Press (2006).

[4] M.J.M. Hill, *On a spherical voltex*, Philos. Trans. R, Soc. London, Ser. A **185**, 213-245 (1894).

[5] A.W. Marris and M.G. Aswani, *On the general impossibility of controllable axisymmetric Navier-Stokes motions*, Arch. Rat. Mech. Anal. **63**, 107-153 (1977).

[6] V. O'Brien, *Steady spheroidal vortices – more exact solutions to the Navier-Stokes equation*, Quart. Appl. Math. **19**, 163-168 (1961).

[7] R.M. Terrill, *An exact solution for flow in a porous pipe*, Z. Angew. Math. Phys. **33**, 547-552 (1982).

[8] C.Y. Wang, *Exact solution of the Navier-Stokes equations – the generalized Beltrami flows, review and extension*, Acta Mech. **81**, 69-74 (1990).

[9] C.Y. Wang, *Exact solution of the steady-state Navier-Stokes equations*, Annu. Rev. Fluid Mech. **23**, 159-177 (1991).

[10] S. Weinbaum and V. O'Brien, *Exact Navier-Stokes solutions including swirl and cross flow*, Phys. Fluids **10**, 1438-1447 (1967).